

A version of Cartan's Theorem A for coherent sheaves on real affine varieties

Tomasz Kowalczyk

Jagiellonian University in Kraków

Łódź, January 12, 2018

Theorem 1 (Serre)

Let X be an affine variety over an algebraically closed field and \mathcal{F} be a coherent sheaf on X . Then \mathcal{F} is spanned by its global sections.

Theorem 1 (Serre)

Let X be an affine variety over an algebraically closed field and \mathcal{F} be a coherent sheaf on X . Then \mathcal{F} is spanned by its global sections.

X will be a non-singular real affine variety with structure sheaf \mathcal{O}_X .

Definition 2

A sheaf \mathcal{F} of \mathcal{O}_X -Modules is called coherent if there exists a finite Zariski open covering $\{U_i\}_{i=1}^n$ of X such that for every U_i there is an exact sequence of sheaves

$$\mathcal{O}_X^{p_i}|_{U_i} \xrightarrow{\phi_i} \mathcal{O}_X^{q_i}|_{U_i} \xrightarrow{\psi_i} \mathcal{F}|_{U_i} \rightarrow 0.$$

There are several cases when Theorem A is true:

There are several cases when Theorem A is true:

- 1 complex analytic geometry,

There are several cases when Theorem A is true:

- ① complex analytic geometry,
- ② algebraic geometry over algebraically closed field (Serre)

There are several cases when Theorem A is true:

- 1 complex analytic geometry,
- 2 algebraic geometry over algebraically closed field (Serre)
- 3 scheme theory (Grothendieck, Hartshorne)

There are several cases when Theorem A is true:

- 1 complex analytic geometry,
- 2 algebraic geometry over algebraically closed field (Serre)
- 3 scheme theory (Grothendieck, Hartshorne)
- 4 regular geometry (Fichou, Huisman, Mangolte, Monnier)

There are several cases when Theorem A is true:

- 1 complex analytic geometry,
- 2 algebraic geometry over algebraically closed field (Serre)
- 3 scheme theory (Grothendieck, Hartshorne)
- 4 regular geometry (Fichou, Huisman, Mangolte, Monnier)
- 5 regular geometry over Henselian valued fields (Nowak)

Example 1

Let $P = X^2(X - 1)^2 + Y^2 \in \mathbb{R}[X, Y]$. The polynomial P is irreducible and has only two zeros $c_1 = (0, 0)$ and $c_2 = (1, 0)$ in \mathbb{R}^2 . Put $U_i = \mathbb{R}^2 \setminus \{c_i\}$. The transition function

$$g_{1,2} : U_1 \cap U_2 \rightarrow GL(1, \mathbb{R}) = \mathbb{R}^*$$

$$(x, y) \mapsto P(x, y)$$

defines a vector bundle of rank 1 over \mathbb{R}^2 .

Example 1

Let $P = X^2(X - 1)^2 + Y^2 \in \mathbb{R}[X, Y]$. The polynomial P is irreducible and has only two zeros $c_1 = (0, 0)$ and $c_2 = (1, 0)$ in \mathbb{R}^2 . Put $U_i = \mathbb{R}^2 \setminus \{c_i\}$. The transition function

$$g_{1,2} : U_1 \cap U_2 \rightarrow GL(1, \mathbb{R}) = \mathbb{R}^*$$

$$(x, y) \mapsto P(x, y)$$

defines a vector bundle of rank 1 over \mathbb{R}^2 . Global sections can be described as a pair (s_1, s_2) where $s_i : U_i \rightarrow \mathbb{R}$ are regular functions and $s_1 = g_{1,2}s_2$. Set $s_i = \frac{f_i}{h_i}$ where f_i, h_i are relatively prime polynomials. Then $f_1h_2 = Pf_2h_1$. Since P does not divide h_2 we obtain that $f_1 = \lambda Pf_2$ and $h_2 = \lambda^{-1}h_1$, where $\lambda \in \mathbb{R}^*$. This shows that every algebraic global section of this bundle vanishes at c_2 .

By a multi-blowup we mean a finite composition of blowups along smooth centers. The basic tool used in these proofs is transformation to simple normal crossing.

By a multi-blowup we mean a finite composition of blowups along smooth centers. The basic tool used in these proofs is transformation to simple normal crossing.

Definition 3

We say that a regular function $g : X \rightarrow \mathbb{R}$ on a non-singular real algebraic variety of dimension d is a simple normal crossing if in a neighbourhood of each point $a \in X$ one has

$$g(x) = u(x)x^\alpha = u(x)x_1^{\alpha_1}x_2^{\alpha_2} \dots x_d^{\alpha_d}$$

where $u(x)$ is a unit at a , $\alpha \in \mathbb{N}^d$ and $x = (x_1, x_2, \dots, x_d)$ are local coordinates near a , i.e. $x_1, x_2, \dots, x_d \in \mathcal{O}_{a,X}$ is a regular system of parameters of the local ring $\mathcal{O}_{a,X}$.

By a multi-blowup we mean a finite composition of blowups along smooth centers. The basic tool used in these proofs is transformation to simple normal crossing.

Definition 3

We say that a regular function $g : X \rightarrow \mathbb{R}$ on a non-singular real algebraic variety of dimension d is a simple normal crossing if in a neighbourhood of each point $a \in X$ one has

$$g(x) = u(x)x^\alpha = u(x)x_1^{\alpha_1}x_2^{\alpha_2} \dots x_d^{\alpha_d}$$

where $u(x)$ is a unit at a , $\alpha \in \mathbb{N}^d$ and $x = (x_1, x_2, \dots, x_d)$ are local coordinates near a , i.e. $x_1, x_2, \dots, x_d \in \mathcal{O}_{a,X}$ is a regular system of parameters of the local ring $\mathcal{O}_{a,X}$.

Theorem 4

*Let f_1, f_2, \dots, f_k be regular functions on X . Then there exists a multi-blowup $\sigma : \tilde{X} \rightarrow X$ such that $\sigma^*f_1, \sigma^*f_2, \dots, \sigma^*f_k$ are simple normal crossings.*

Lemma 5

Let X be a non-singular real affine variety and $U = X \setminus \{Q = 0\}$ a Zariski open subset of X . Every regular function f on U can be written in the form $f = \frac{g}{P}$ where g, P are global regular functions on X and $V(P) \subset V(Q)$.

Lemma 5

Let X be a non-singular real affine variety and $U = X \setminus \{Q = 0\}$ a Zariski open subset of X . Every regular function f on U can be written in the form $f = \frac{g}{P}$ where g, P are global regular functions on X and $V(P) \subset V(Q)$.

Lemma 6

Let X be a non-singular real affine variety, Q a regular function on X and $U := X \setminus \{Q = 0\}$. Then for any $f \in \mathcal{O}_X(U)$ there exists a multi-blowup $\sigma : \tilde{X} \rightarrow X$ and a positive integer N such that $(Q^N f)^\sigma$ can be extended to a global regular function, i.e. $(Q^N f)^\sigma \in \mathcal{O}_{\tilde{X}}(\tilde{X})$.

If the function f is of the form $f = \frac{g}{P}$ as a consequence of proof we get that

$$(Q^N)^\sigma \in P^\sigma \mathcal{O}_{\tilde{X}}(\tilde{X})$$

We will now prove a crucial lemma

We will now prove a crucial lemma

Lemma 7

Let X be a non-singular real affine variety, \mathcal{F} a coherent sheaf on X . For any $Q \in \mathcal{O}_X(X)$ and a section $s \in \mathcal{F}(X)$ such that $s|_U = 0$ with $U = X \setminus \{Q = 0\}$, there exists a multi-blowup $\sigma : \tilde{X} \rightarrow X$ and a positive integer N such that $(Q^N)^\sigma \sigma^ s = 0$ in $\sigma^* \mathcal{F}(\tilde{X})$.*

Proof of Lemma 7

By quasi-compactness, there is a finite Zariski open covering $\{U_i\}_{i=1}^n$ of X such that for each i we have:

Proof of Lemma 7

By quasi-compactness, there is a finite Zariski open covering $\{U_i\}_{i=1}^n$ of X such that for each i we have:

1) a presentation

$$\mathcal{O}_X^p|_{U_i} \xrightarrow{\phi_i} \mathcal{O}_X^q|_{U_i} \xrightarrow{\psi_i} \mathcal{F}|_{U_i} \rightarrow 0.$$

Proof of Lemma 7

By quasi-compactness, there is a finite Zariski open covering $\{U_i\}_{i=1}^n$ of X such that for each i we have:

1) a presentation

$$\mathcal{O}_X^p|_{U_i} \xrightarrow{\phi_i} \mathcal{O}_X^{q_i}|_{U_i} \xrightarrow{\psi_i} \mathcal{F}|_{U_i} \rightarrow 0.$$

2) $s|_{U_i} = \psi_i(t_i)$ for some $t_i \in \mathcal{O}_X^{q_i}(U_i)$.

Proof of Lemma 7

By quasi-compactness, there is a finite Zariski open covering $\{U_i\}_{i=1}^n$ of X such that for each i we have:

1) a presentation

$$\mathcal{O}_X^p|_{U_i} \xrightarrow{\phi_i} \mathcal{O}_X^{q_i}|_{U_i} \xrightarrow{\psi_i} \mathcal{F}|_{U_i} \rightarrow 0.$$

2) $s|_{U_i} = \psi_i(t_i)$ for some $t_i \in \mathcal{O}_X^{q_i}(U_i)$.

By the assumption, $t_{ix} \in \phi_i(\mathcal{O}_{x,X}^p)$ for each $x \in U_i \cap U$, and thus

$$t_{ix} = \sum_{j=1}^p f_{ijx} \phi_i(e_j)_x,$$

where $e_j = (0, 0, \dots, 0, \underset{j}{1}, 0, \dots, 0) \in \mathcal{O}_X^p(X)$ and $f_{ijx} \in \mathcal{O}_{x,X}$.

Proof of Lemma 7

Define the ideal

$$I_i = \{P_i \in \mathcal{O}_X(X) : P_i t_i \in \phi_i(\mathcal{O}_X^p)(U_i)\} \quad i = 1, 2, \dots, n.$$

Claim: $V(I_i) \cap U_i \subset V(Q) \cap U_i$.

Proof of Lemma 7

Define the ideal

$$I_i = \{P_i \in \mathcal{O}_X(X) : P_i t_i \in \phi_i(\mathcal{O}_X^p)(U_i)\} \quad i = 1, 2, \dots, n.$$

Claim: $V(I_i) \cap U_i \subset V(Q) \cap U_i$.

Proof.

If $x \in U_i \setminus V(Q)$, then

$$f_{ijx} = \frac{g_{ijx}}{h_{ijx}} \quad \text{where } g_{ijx}, h_{ijx} \in \mathcal{O}_X(X), h_{ijx}(x) \neq 0.$$

Define $P_i := \prod_{j=1}^p h_{ijx}$. Then

$$P_i t_i \in \sum_{j=1}^p \mathcal{O}_X(X) \phi_i(e_j|_{U_i}),$$

and thus $P_i \in I_i$. Since $P_i(x) \neq 0$ we get, $x \notin V(I_i)$. □

Proof of Lemma 7

Let $P_{i1}, P_{i2}, \dots, P_{ir_i}$ be generators of I_i . Taking $P_i := P_{i1}^2 + \dots + P_{ir_i}^2 \in I_i$, we get $V(P_i) \subset V(Q)$.

Proof of Lemma 7

Let $P_{i1}, P_{i2}, \dots, P_{ir_i}$ be generators of I_i . Taking $P_i := P_{i1}^2 + \dots + P_{ir_i}^2 \in I_i$, we get $V(P_i) \subset V(Q)$.

Consider the case $i = 1$.

Proof of Lemma 7

Let $P_{i1}, P_{i2}, \dots, P_{ir_i}$ be generators of I_i . Taking $P_i := P_{i1}^2 + \dots + P_{ir_i}^2 \in I_i$, we get $V(P_i) \subset V(Q)$.

Consider the case $i = 1$. By the reasoning as in the proof of Lemma 6, there exist a multi-blowup $\sigma_1 : X_1 \rightarrow X$ and a positive integer N_1 such that

$$(Q^{N_1})^{\sigma_1} \in P_1^{\sigma_1} \mathcal{O}_{X_1}(X_1).$$

Proof of Lemma 7

Let $P_{i1}, P_{i2}, \dots, P_{ir_i}$ be generators of I_i . Taking $P_i := P_{i1}^2 + \dots + P_{ir_i}^2 \in I_i$, we get $V(P_i) \subset V(Q)$.

Consider the case $i = 1$. By the reasoning as in the proof of Lemma 6, there exist a multi-blowup $\sigma_1 : X_1 \rightarrow X$ and a positive integer N_1 such that

$$(Q^{N_1})^{\sigma_1} \in P_1^{\sigma_1} \mathcal{O}_{X_1}(X_1).$$

Hence

$$(Q^{N_1})^{\sigma_1} \sigma_1^* t_1 \in \phi_1^{\sigma_1}(\mathcal{O}_{X_1}^p)(U_1^{\sigma_1})$$

Proof of Lemma 7

Let $P_{i1}, P_{i2}, \dots, P_{ir_i}$ be generators of I_i . Taking $P_i := P_{i1}^2 + \dots + P_{ir_i}^2 \in I_i$, we get $V(P_i) \subset V(Q)$.

Consider the case $i = 1$. By the reasoning as in the proof of Lemma 6, there exist a multi-blowup $\sigma_1 : X_1 \rightarrow X$ and a positive integer N_1 such that

$$(Q^{N_1})^{\sigma_1} \in P_1^{\sigma_1} \mathcal{O}_{X_1}(X_1).$$

Hence

$$(Q^{N_1})^{\sigma_1} \sigma_1^* t_1 \in \phi_1^{\sigma_1}(\mathcal{O}_{X_1}^p)(U_1^{\sigma_1})$$

and thus

$$(Q^{N_1})^{\sigma_1} \sigma_1^* s|_{U_1^{\sigma_1}} = 0.$$

Proof of Lemma 7

Note that if $\sigma_1 : X_1 \rightarrow X$ is a blowup and if a covering $\{U_i\}_{i=1}^n$ of X satisfies conditions 1) and 2), then so does the covering $\{U_i^{\sigma_1}\}_{i=1}^n$.

Proof of Lemma 7

Note that if $\sigma_1 : X_1 \rightarrow X$ is a blowup and if a covering $\{U_i\}_{i=1}^n$ of X satisfies conditions 1) and 2), then so does the covering $\{U_i^{\sigma_1}\}_{i=1}^n$. Now we can repeat the reasoning for $U_2^{\sigma_1}$ to obtain a positive integer $N_2 \geq N_1$ and a multi-blowup $\sigma_2 : X_2 \rightarrow X_1$ such that

$$(Q^{N_2})^{\sigma_1 \circ \sigma_2} (\sigma_1 \circ \sigma_2)^* s|_{(U_1 \cup U_2)^{\sigma_1 \circ \sigma_2}} = 0$$

and so on.

Proof of Lemma 7

Note that if $\sigma_1 : X_1 \rightarrow X$ is a blowup and if a covering $\{U_i\}_{i=1}^n$ of X satisfies conditions 1) and 2), then so does the covering $\{U_i^{\sigma_1}\}_{i=1}^n$. Now we can repeat the reasoning for $U_2^{\sigma_1}$ to obtain a positive integer $N_2 \geq N_1$ and a multi-blowup $\sigma_2 : X_2 \rightarrow X_1$ such that

$$(Q^{N_2})^{\sigma_1 \circ \sigma_2} (\sigma_1 \circ \sigma_2)^* s|_{(U_1 \cup U_2)^{\sigma_1 \circ \sigma_2}} = 0$$

and so on. We continue this process and obtain a positive integer $N := N_n \geq N_{n-1} \geq \dots \geq N_1$ and a multi-blowup

$\sigma := \sigma_1 \circ \dots \circ \sigma_n : \tilde{X} \rightarrow X$, $\tilde{X} := X_n$, such that $(Q^N)^\sigma \sigma^* s$ vanishes on $\tilde{X} = (U_1 \cup U_2 \cup \dots \cup U_n)^\sigma$. This finishes the proof.

Let $f : X \rightarrow Y$ be a morphism of real algebraic varieties.

Lemma 8

If \mathcal{G} is of finite type or coherent sheaf of \mathcal{O}_Y -Modules generated by sections $s_1, s_2, \dots, s_k \in \mathcal{G}(Y)$, then the pull-back $f^\mathcal{G}$ is generated by the pull-back $f^*s_1, f^*s_2, \dots, f^*s_k \in (f^*\mathcal{G})(X)$.*

Lemma 9

Let \mathcal{F} be a coherent sheaf on a non-singular real affine variety with local presentations

$$\mathcal{O}_X^p|_{U_i} \xrightarrow{\phi_i} \mathcal{O}_X^{q_i}|_{U_i} \xrightarrow{\psi_i} \mathcal{F}|_{U_i} \rightarrow 0 \quad i = 1, 2, \dots, n$$

on a finite Zariski open covering $\{U_1, U_2, \dots, U_n\}$. Consider a finite family of Zariski open sets

$$V_j = X \setminus \{Q_j = 0\}, \quad j = 1, 2, \dots, m$$

where Q_j are regular functions on X , and sections $s_j \in \mathcal{F}(V_j)$. Assume that every V_j is contained in $U_{i(j)}$ for some $i(j) = 1, 2, \dots, n$ and that for each s_j there is a section $t_j \in \mathcal{O}_X^{q_{i(j)}}(V_j)$ such that $\psi_{i(j)}(V_j)(t_j) = s_j$. Then there exists a positive integer N and a multi-blowup $\sigma : \tilde{X} \rightarrow X$ such that every section $(Q_j^N)^\sigma \sigma^* s_j$ $j = 1, 2, \dots, m$ extends to a global section on \tilde{X} .

Proof of Lemma 9

Since taking pull-back under a multi-blowup does not affect the assumptions, it suffices to consider only one $j = 1, 2, \dots, m$.

Proof of Lemma 9

Since taking pull-back under a multi-blowup does not affect the assumptions, it suffices to consider only one $j = 1, 2, \dots, m$. So fix a j and let $t_{ji} = t_j|_{U_i \cap V_j} = (t_{ji1}, t_{ji2}, \dots, t_{jiqu_i})$ where

$$t_{jil} = \frac{t_{jil1}}{t_{jil2}}, \quad t_{jil1}, t_{jil2} \in \mathcal{O}_X(X)$$

and

$$V(t_{jil2}) \cap U_i \subset V(Q_j) \cap U_i.$$

Proof of Lemma 9

Since taking pull-back under a multi-blowup does not affect the assumptions, it suffices to consider only one $j = 1, 2, \dots, m$. So fix a j and let $t_{ji} = t_j|_{U_i \cap V_j} = (t_{ji1}, t_{ji2}, \dots, t_{jil}, \dots, t_{jiq_i})$ where

$$t_{jil} = \frac{t_{jil1}}{t_{jil2}}, \quad t_{jil1}, t_{jil2} \in \mathcal{O}_X(X)$$

and

$$V(t_{jil2}) \cap U_i \subset V(Q_j) \cap U_i.$$

Using Lemma 6 we can find a positive integer N_1 and a multi-blowup $\tau_1 : X_1 \rightarrow X$ such that

$$(t_{jil}(Q_j)^{N_1})^{\tau_1} \in \mathcal{O}_{X_1}(U_i^{\tau_1}) \quad \text{for all } j, i, l.$$

Now define $\tilde{s}_{ji} := \psi_i^{\tau_1}((t_{jil}(Q_j)^{N_1})^{\tau_1})$.

Proof of Lemma 9

Then for any two distinct indices i_0, i_1 we have

$$(\widetilde{s}_{ji_0} - \widetilde{s}_{ji_1})|_{U_{i_0}^{\tau_1} \cap U_{i_1}^{\tau_1}} \in (\tau_1^* \mathcal{F})(U_{i_0}^{\tau_1} \cap U_{i_1}^{\tau_1})$$

and

$$(\widetilde{s}_{ji_0} - \widetilde{s}_{ji_1})|_{U_{i_0}^{\tau_1} \cap U_{i_1}^{\tau_1} \cap V_j^{\tau_1}} = 0.$$

Proof of Lemma 9

Then for any two distinct indices i_0, i_1 we have

$$(\widetilde{s}_{ji_0} - \widetilde{s}_{ji_1})|_{U_{i_0}^{\tau_1} \cap U_{i_1}^{\tau_1}} \in (\tau_1^* \mathcal{F})(U_{i_0}^{\tau_1} \cap U_{i_1}^{\tau_1})$$

and

$$(\widetilde{s}_{ji_0} - \widetilde{s}_{ji_1})|_{U_{i_0}^{\tau_1} \cap U_{i_1}^{\tau_1} \cap V_j^{\tau_1}} = 0.$$

By Lemma 7 we can find a multi-blowup $\tau_2 : \widetilde{X} \rightarrow X_1$ and a positive integer N_2 such that

$$(\tau_2^* \widetilde{s}_{ji_0} (Q_j^{N_1+N_2})^{\tau_1 \circ \tau_2} - \tau_2^* \widetilde{s}_{ji_1} (Q_j^{N_1+N_2})^{\tau_1 \circ \tau_2})|_{U_{i_0}^{\tau_1 \circ \tau_2} \cap U_{i_1}^{\tau_1 \circ \tau_2}} = 0.$$

Proof of Lemma 9

Then for any two distinct indices i_0, i_1 we have

$$(\widetilde{s}_{j_{i_0}} - \widetilde{s}_{j_{i_1}})|_{U_{i_0}^{\tau_1} \cap U_{i_1}^{\tau_1}} \in (\tau_1^* \mathcal{F})(U_{i_0}^{\tau_1} \cap U_{i_1}^{\tau_1})$$

and

$$(\widetilde{s}_{j_{i_0}} - \widetilde{s}_{j_{i_1}})|_{U_{i_0}^{\tau_1} \cap U_{i_1}^{\tau_1} \cap V_j^{\tau_1}} = 0.$$

By Lemma 7 we can find a multi-blowup $\tau_2 : \widetilde{X} \rightarrow X_1$ and a positive integer N_2 such that

$$(\tau_2^* \widetilde{s}_{j_{i_0}}(Q_j^{N_1+N_2})^{\tau_1 \circ \tau_2} - \tau_2^* \widetilde{s}_{j_{i_1}}(Q_j^{N_1+N_2})^{\tau_1 \circ \tau_2})|_{U_{i_0}^{\tau_1 \circ \tau_2} \cap U_{i_1}^{\tau_1 \circ \tau_2}} = 0.$$

Considering all distinct pairs of indices, we can assume that the differences as above vanish for all those pairs. Therefore the sections

$$(\tau_2^* \widetilde{s}_{j_i}(Q_j^{N_1+N_2})^{\tau_1 \circ \tau_2})|_{U_i^{\tau_1 \circ \tau_2}}, \quad i = 1, 2, \dots, n,$$

glue together to a global section on \widetilde{X} . Thus $\sigma := \tau_1 \circ \tau_2 : \widetilde{X} \rightarrow X$ is the multi-blowup we are looking for.

Lemma 10

Let \mathcal{F} be a sheaf of \mathcal{O}_X -Modules of finite type and let $s_1, s_2, \dots, s_k \in \mathcal{F}(U)$ be sections of \mathcal{F} on a neighbourhood U of a point $a \in X$. If $s_{1a}, s_{2a}, \dots, s_{ka}$ generate \mathcal{F}_a , then $s_{1x}, s_{2x}, \dots, s_{kx}$ generate \mathcal{F}_x for all x sufficiently close to a .

Lemma 10

Let \mathcal{F} be a sheaf of \mathcal{O}_X -Modules of finite type and let $s_1, s_2, \dots, s_k \in \mathcal{F}(U)$ be sections of \mathcal{F} on a neighbourhood U of a point $a \in X$. If $s_{1a}, s_{2a}, \dots, s_{ka}$ generate \mathcal{F}_a , then $s_{1x}, s_{2x}, \dots, s_{kx}$ generate \mathcal{F}_x for all x sufficiently close to a .

Assume that, under the above assumptions, $U := X \setminus \{Q = 0\}$ with some $Q \in \mathcal{O}_X(X)$. Then the sections $Q^n s_1, Q^n s_2, \dots, Q^n s_k$ generate every stalk sufficiently close to a because the function Q is invertible in $\mathcal{O}_{a,X}$ for every $a \in U$. Now we are ready to prove the main theorem.

Lemma 10

Let \mathcal{F} be a sheaf of \mathcal{O}_X -Modules of finite type and let $s_1, s_2, \dots, s_k \in \mathcal{F}(U)$ be sections of \mathcal{F} on a neighbourhood U of a point $a \in X$. If $s_{1a}, s_{2a}, \dots, s_{ka}$ generate \mathcal{F}_a , then $s_{1x}, s_{2x}, \dots, s_{kx}$ generate \mathcal{F}_x for all x sufficiently close to a .

Assume that, under the above assumptions, $U := X \setminus \{Q = 0\}$ with some $Q \in \mathcal{O}_X(X)$. Then the sections $Q^n s_1, Q^n s_2, \dots, Q^n s_k$ generate every stalk sufficiently close to a because the function Q is invertible in $\mathcal{O}_{a,X}$ for every $a \in U$. Now we are ready to prove the main theorem.

Theorem 11

Let \mathcal{F} be a coherent sheaf on a non-singular real affine variety X . Then there exist a multi-blowup $\sigma : \tilde{X} \rightarrow X$ and finitely many global sections s_1, s_2, \dots, s_k on \tilde{X} which generate every stalk $(\sigma^* \mathcal{F})_y$, $y \in \tilde{X}$.

Proof of Theorem 11

Consider a finite Zariski open covering $\{U_1, U_2, \dots, U_n\}$ of X with local presentation of the sheaf \mathcal{F}

$$\mathcal{O}_X^p|_{U_i} \xrightarrow{\phi_i} \mathcal{O}_X^{q_i}|_{U_i} \xrightarrow{\psi_i} \mathcal{F}|_{U_i} \rightarrow 0.$$

Proof of Theorem 11

Consider a finite Zariski open covering $\{U_1, U_2, \dots, U_n\}$ of X with local presentation of the sheaf \mathcal{F}

$$\mathcal{O}_X^p|_{U_i} \xrightarrow{\phi_i} \mathcal{O}_X^{q_i}|_{U_i} \xrightarrow{\psi_i} \mathcal{F}|_{U_i} \rightarrow 0.$$

By Lemma 11, for any point $a \in X$ there are finitely many sections

$$s_{a1}, s_{a2}, \dots, s_{am_a} \in \mathcal{F}(V_a), \quad m_a \in \mathbb{N},$$

on a Zariski open neighbourhood V_a of a , contained in U_i for some $i = 1, 2, \dots, n$, which generate \mathcal{F} over V_a .

Proof of Theorem 11

Consider a finite Zariski open covering $\{U_1, U_2, \dots, U_n\}$ of X with local presentation of the sheaf \mathcal{F}

$$\mathcal{O}_X^p|_{U_i} \xrightarrow{\phi_i} \mathcal{O}_X^{q_i}|_{U_i} \xrightarrow{\psi_i} \mathcal{F}|_{U_i} \rightarrow 0.$$

By Lemma 11, for any point $a \in X$ there are finitely many sections

$$s_{a1}, s_{a2}, \dots, s_{am_a} \in \mathcal{F}(V_a), \quad m_a \in \mathbb{N},$$

on a Zariski open neighbourhood V_a of a , contained in U_i for some $i = 1, 2, \dots, n$, which generate \mathcal{F} over V_a . After shrinking V_a , we can also assume that $s_{ak} = \psi_i(t_{ak})$ for some $t_{ak} \in \mathcal{O}_X^{q_i}(V_a)$, $k = 1, 2, \dots, m_a$.

Proof of Theorem 11

Consider a finite Zariski open covering $\{U_1, U_2, \dots, U_n\}$ of X with local presentation of the sheaf \mathcal{F}

$$\mathcal{O}_X^p|_{U_i} \xrightarrow{\phi_i} \mathcal{O}_X^{q_i}|_{U_i} \xrightarrow{\psi_i} \mathcal{F}|_{U_i} \rightarrow 0.$$

By Lemma 11, for any point $a \in X$ there are finitely many sections

$$s_{a1}, s_{a2}, \dots, s_{am_a} \in \mathcal{F}(V_a), \quad m_a \in \mathbb{N},$$

on a Zariski open neighbourhood V_a of a , contained in U_i for some $i = 1, 2, \dots, n$, which generate \mathcal{F} over V_a . After shrinking V_a , we can also assume that $s_{ak} = \psi_i(t_{ak})$ for some $t_{ak} \in \mathcal{O}_X^{q_i}(V_a)$, $k = 1, 2, \dots, m_a$. By quasi-compactness, we can find a finite covering $V_j := V_{a_j}$, $j = 1, 2, \dots, m$ of X . Clearly, every V_j is contained in $U_{i(j)}$ for some $i(j) = 1, 2, \dots, n$.

Proof of Theorem 11

Consider a finite Zariski open covering $\{U_1, U_2, \dots, U_n\}$ of X with local presentation of the sheaf \mathcal{F}

$$\mathcal{O}_X^p|_{U_i} \xrightarrow{\phi_i} \mathcal{O}_X^{q_i}|_{U_i} \xrightarrow{\psi_i} \mathcal{F}|_{U_i} \rightarrow 0.$$

By Lemma 11, for any point $a \in X$ there are finitely many sections

$$s_{a1}, s_{a2}, \dots, s_{am_a} \in \mathcal{F}(V_a), \quad m_a \in \mathbb{N},$$

on a Zariski open neighbourhood V_a of a , contained in U_i for some $i = 1, 2, \dots, n$, which generate \mathcal{F} over V_a . After shrinking V_a , we can also assume that $s_{ak} = \psi_i(t_{ak})$ for some $t_{ak} \in \mathcal{O}_X^{q_i}(V_a)$, $k = 1, 2, \dots, m_a$. By quasi-compactness, we can find a finite covering $V_j := V_{a_j}$, $j = 1, 2, \dots, m$ of X . Clearly, every V_j is contained in $U_{i(j)}$ for some $i(j) = 1, 2, \dots, n$. Put

$$s_{jk} = s_{a_j k} \quad \text{and} \quad t_{jk} = t_{a_j k}$$

for $j = 1, 2, \dots, m$, $k = 1, 2, \dots, m_j = m_{a_j}$.

Proof of Theorem 11

Then $s_{jk} = \psi_{i(j)}(t_{jk})$ and the sections s_{jk} , $k = 1, 2, \dots, m_j$ generate \mathcal{F} over V_j .

Proof of Theorem 11

Then $s_{jk} = \psi_{i(j)}(t_{jk})$ and the sections s_{jk} , $k = 1, 2, \dots, m_j$ generate \mathcal{F} over V_j . The sets $X \setminus V_j$ are Zariski closed and thus are of the form

$$X \setminus V_j = \{Q_j = 0\}, \quad j = 1, 2, \dots, m$$

for some regular functions Q_j on X .

Proof of Theorem 11

Then $s_{jk} = \psi_{i(j)}(t_{jk})$ and the sections s_{jk} , $k = 1, 2, \dots, m_j$ generate \mathcal{F} over V_j . The sets $X \setminus V_j$ are Zariski closed and thus are of the form

$$X \setminus V_j = \{Q_j = 0\}, \quad j = 1, 2, \dots, m$$

for some regular functions Q_j on X . It follows from Lemma 9 that there exist a multi-blowup $\sigma : \tilde{X} \rightarrow X$ and a positive integer N such that for each $j = 1, 2, \dots, m$ the sections

$$(Q_j^N)^\sigma \sigma^* s_{jk} \quad k = 1, 2, \dots, m_j,$$

extends to global sections $\widetilde{s}_{jk} \in \sigma^* \mathcal{F}(\tilde{X})$. Since $\{\sigma^{-1}(V_j)\}_{j=1}^m$ is a Zariski open covering of \tilde{X} , it is easy to check that the global sections

$$\widetilde{s}_{jk}, \quad j = 1, 2, \dots, m, \quad k = 1, 2, \dots, m_j$$

generate the pull-back $(\sigma^* \mathcal{F})_y$ for every $y \in \tilde{X}$. This finishes the proof.

Corollary 1

Let \mathcal{F} be a coherent sheaf on a non-singular real affine variety X . Then there exists a multi-blowup $\sigma : \tilde{X} \rightarrow X$ such that the pull-back $\sigma^*\mathcal{F}$ admits a global presentation:

$$\mathcal{O}_{\tilde{X}}^p \rightarrow \mathcal{O}_{\tilde{X}}^q \rightarrow \sigma^*\mathcal{F} \rightarrow 0$$

Corollary 1

Let \mathcal{F} be a coherent sheaf on a non-singular real affine variety X . Then there exists a multi-blowup $\sigma : \tilde{X} \rightarrow X$ such that the pull-back $\sigma^*\mathcal{F}$ admits a global presentation:

$$\mathcal{O}_{\tilde{X}}^p \rightarrow \mathcal{O}_{\tilde{X}}^q \rightarrow \sigma^*\mathcal{F} \rightarrow 0$$

Connections with works of Tognoli:

Corollary 2

Let \mathcal{F} be a coherent sheaf on a non-singular real affine variety X . Then there exists a multi-blowup $\sigma : \tilde{X} \rightarrow X$ such that $\sigma^*\mathcal{F}$ is an A -coherent sheaf.